

Hodge-Frobenius equations and the Hodge-Bäcklund transformation

Antonella Marini* and Thomas H. Otway†

*Department of Mathematics, Yeshiva University,
New York, New York 10033*

Abstract

Linear and nonlinear Hodge-like systems for 1-forms are studied, with an assumption equivalent to complete integrability substituted for the requirement of closure under exterior differentiation. The systems are placed in a variational context and properties of critical points investigated. Certain standard choices of energy density are related by Bäcklund transformations which employ basic properties of the Hodge involution. These *Hodge-Bäcklund transformations* yield invariant forms of classical Bäcklund transformations that arise in diverse contexts. Some extensions to higher-degree forms are indicated. *MSC2000*: 35A15, 58J72, 58A14.

Key words: nonlinear Hodge theory, Bäcklund transformation, Frobenius Theorem

1 Introduction

The study of vectors which are both divergence-free and curl-free can be traced back at least to Helmholtz's analysis of vortices and gradients [14]. The generalization to differential forms which are both closed and co-closed under exterior differentiation is the content of the Hodge equations; see, *e.g.*, [25], Ch. 7. The divergence-free condition is frequently relaxed in variational contexts, but generalizations of the curl-free condition remain rather rare.

Our goal is to study both linear and nonlinear variants of the Hodge equations for differential forms which are neither co-closed nor closed, but which satisfy milder conditions having physical and geometric significance.

*Permanent address: Dipartimento di Matematica, Università di L'Aquila, 67100 L'Aquila, Italy; email: marini@dm.univaq.it

†email: otway@yu.edu

1.1 Organization of the paper

Sections 1-4 are mainly expository. We introduce the topic in Sec. 1.2 with an example from fluid dynamics. Section 2 presents the equations in an invariant context. The linear case is studied in Sec. 3, largely as motivation for the considerably more complex nonlinear case. Two geometric analogies are discussed in Sec. 4. Technical results on the properties of solutions are presented in Sec. 5. Section 6 shows that ideas introduced, in very different contexts, by Yang [43] and by Magnanini and Talenti [19]-[21], can be given a unified interpretation in terms of equations studied in the preceding sections.

The proofs of Theorems 5, 6, and 8, which are based on rather straightforward applications of nonlinear elliptic theory, are collected in an appendix, Sec. 7. We note that these applications are only straightforward once solutions have been associated to a uniformly sub-elliptic operator; this is accomplished in Sec. 5.2. The methods used in that section to derive uniform estimates are elementary and, in particular, do not require a delicate limiting argument that has become known as *Shiffman regularization* (*c.f.* [34] and the appendix by R. J. Sibner to [35]).

1.2 A motivating example: steady, ideal flow

In models for the steady, adiabatic and isentropic flow of an ideal fluid, conservation of mass is represented by the *continuity equation*

$$\nabla \cdot (\rho (|\mathbf{v}|^2) \mathbf{v}) = 0, \quad (1)$$

where $\mathbf{v}$ denotes flow velocity and ρ denotes mass density. The dependence of ρ on $|\mathbf{v}|^2$ is a consequence of compressibility; in the incompressible limit, eq. (1) says merely that the vector $\mathbf{v}$ has zero divergence. If the fluid is irrotational, then the velocity is curl-free in the sense that

$$\nabla \times \mathbf{v} = 0. \quad (2)$$

Condition (2) implies, by the Poincaré Lemma, that there exists locally a scalar flow potential $\varphi(\mathbf{x})$, where $\mathbf{x} \in \mathbb{R}^3$ denotes the position of a particle in the flow.

Perhaps the mildest weakening of the irrotationality condition results from replacing (2) by the integrability condition

$$\mathbf{v} \cdot \nabla \times \mathbf{v} = 0. \quad (3)$$

The replacement of the linear condition (2) by the nonlinear condition (3) as a side condition to eq. (1) is likely to result in singular solutions, even in the subsonic regime. The usual arguments for reducing ρ to the conventional form, which depend on smoothness (see, *e.g.*, Ch. 1.2 of [4]), would not necessarily apply in such cases. This suggests that we consider whether a useful *a priori* bound can be placed on the size of the singular set for solutions of systems having the general form (1), (3). In Sec. 5.2 we take the first step toward an answer to this difficult question, deriving sufficient conditions under which a

solution remains bounded on an apparent singular set of given codimension. Despite the physical motivation (here and in various other examples scattered throughout the text), our main interest in this paper is in deriving hypotheses which are mathematically natural and apply to large classes of mass densities.

2 An invariant formulation

Thus we generalize the mathematical context of Sec. 1.2. Let Ω be an open, finite domain of $\mathbb{R}^n$, $n \geq 2$, satisfying an interior sphere condition. Consider the system ([26], Sec. VI; [27], Sec. 4)

$$\delta(\rho(Q)\omega) = 0, \quad (4)$$

$$d\omega = \Gamma \wedge \omega, \quad (5)$$

for scalar-valued 1-forms ω and Γ , where Γ is given and ω is unknown; d is the flat exterior derivative with formal adjoint δ ; Q is a quadratic form in ω given by

$$Q(\omega) = *(\omega \wedge *\omega) \equiv \langle \omega, \omega \rangle, \quad (6)$$

where $*$: $\Lambda^k \rightarrow \Lambda^{n-k}$ is the Hodge involution; ρ is a positive, continuously differentiable function of Q (but a possibly singular function of $\mathbf{x}$).

Using (5), we find that

$$\omega \wedge d\omega = -\Gamma \wedge \omega \wedge \omega = 0. \quad (7)$$

A 1-form ω that is the pointwise Riemannian inner product with a vector field $\mathbf{v}$ is said to be *dual* to $\mathbf{v}$. In this case the left-hand side of (7) is equivalent to the left-hand side of (3), and any solution of ω of eq. (5) is dual to a solution $\mathbf{v}$ of eq. (3).

The system (4), (5) is uniformly elliptic provided the differential inequality

$$0 < \kappa_1 \leq \frac{(d/dQ)[Q\rho^2(Q)]}{\rho(Q)} \leq \kappa_2 < \infty \quad (8)$$

is satisfied for constants κ_1, κ_2 . In the context of fluid dynamics, one typically encounters the weaker condition

$$0 < \rho^2(Q) + 2Q\rho'(Q)\rho(Q). \quad (9)$$

Ideal flow governed by eq. (1) is subsonic provided (9) is satisfied. Moreover, there is typically a critical value Q_{crit} such that the right-hand side of (9) tends to zero in the limit as Q tends to Q_{crit} . In this case, eqs. (4), (5) with ρ satisfying (9) are elliptic, but not uniformly so, and this condition has mathematical as well as physical interest.

If $\Gamma \equiv 0$, then the system (4), (5) degenerates to the *nonlinear Hodge equations* introduced in [36] on the basis of a conjecture in [3]. In that case condition

(5) generates a cohomology class, which is not true in the more general case studied here.

We show in Sec. 2.1 that whenever ω is a 1-form, eq. (5) possesses solutions of the form

$$\omega = e^\eta du, \quad (10)$$

where u and η are 0-forms, and Γ can be made *exact*: we can write $\Gamma = d\eta$. When ω is a k -form, a more general representation applies; that representation is discussed in Sec. 2.2.

In Sec. 5.1 we introduce a variant of eq. (4) for differential k -forms ($k \geq 1$) satisfying eq. (5) with Γ exact ($\Gamma \equiv d\eta$), namely

$$\delta [\rho(Q)\omega] = (-1)^{n(k+1)} * (d\eta \wedge * \rho(Q)\omega), \quad (11)$$

which arises as a variational equation of the *nonlinear Hodge energy*

$$E = \frac{1}{2} \int_{\Omega} \int_0^Q \rho(s) ds d\Omega. \quad (12)$$

2.1 The Frobenius Theorem

Let Γ be fixed, and define

$$S \equiv \{\omega \in \Lambda(\Omega) \equiv \bigoplus_{k=1}^n \Lambda^k(\Omega) : d\omega = \Gamma \wedge \omega\}.$$

Denote by $I \equiv I(S)$ the ideal generated by S . If $\Gamma \neq 0$, then clearly $dI \neq \{0\}$, so the k -forms $\omega \in S$ do not generate cohomology classes.

Nevertheless, the ideal I is *closed*, i.e., $dI \subset I$. In fact, a differential form $\alpha \in I$ is a linear combination of forms of type $\omega \wedge \beta$ with $\omega \in S$, $\beta \in \Lambda(\Omega)$. The latter satisfy

$$d(\omega \wedge \beta) = \pm \omega \wedge (\Gamma \wedge \beta \pm d\beta),$$

and thus satisfy $d\alpha \in I$. This is an important fact, especially for exterior systems of 1-forms.

Following the approach in Sec. 4-2 of [10], we define an exterior system $\{\omega^a\}$, $a = 1, \dots, r$ of r 1-forms in a space of dimension $n = r + s$ to be *completely integrable* if and only if there exist r independent functions g^a , $a = 1, \dots, r$ such that each of the 1-forms ω^a vanishes on the r -parameter family of s -dimensional hypersurfaces $\{g^a = k^a, a = 1, \dots, r\}$ generated by letting the constants k^a range over all r -tuples of real numbers.

Equivalently, we define $\{\omega^a\}_1^r$ to be *completely integrable* if and only if there exists a nonsingular $r \times r$ matrix of functions ξ_b^a , and r independent functions g^b such that

$$\omega^a = \sum_{b=1}^r \xi_b^a dg^b.$$

The *Frobenius Theorem* asserts that an exterior system $\{\omega^a\}_1^r$ of 1-forms is completely integrable if and only if it generates a closed ideal of $\Lambda(\Omega)$.

Because a 1-form ω satisfying (5) generates a closed ideal, by the Frobenius Theorem it can always be written in the form (10). (In this case, $r = 1$.) Thus

$$d\omega = d\eta \wedge \omega, \quad (13)$$

which shows that Γ can be chosen to be exact. Notice the gauge invariance having the form $\Gamma \rightarrow \tilde{\Gamma} \equiv \Gamma + f(\mathbf{x})\omega$.

For this reason, we call the system (4), (5) the *nonlinear Hodge-Frobenius equations for 1-forms*.

Unfortunately, the Frobenius Theorem does not generalize to forms of arbitrary degree k , as the condition $dI \subset I$ does not imply complete integrability if $k \neq 1$. However, this does not mean that there is nothing to be said about higher-degree forms. Relevant properties of such forms are described in the following section.

2.2 Recursive forms

An exterior differential form of degree k is said to be *recursive with coefficient* Γ if it satisfies (5). Let Ω be star-shaped. It is known that one can define a *homotopy operator* $\mathcal{H} : \Lambda^k(\Omega) \rightarrow \Lambda^{k-1}(\Omega)$, which satisfies

$$\omega = d\mathcal{H}\omega + \mathcal{H}d\omega. \quad (14)$$

This property can be used, among other things, to show that a closed form on a star-shaped domain is exact. We omit the formal definition of this operator (for this and further details see Sec. 5-3 of [10]), and only describe its main properties:

- (a) $\mathcal{H}$ is linear
- (b) $\mathcal{H}^2 = 0$
- (c) $\mathcal{H}d\mathcal{H} = \mathcal{H}$, $d\mathcal{H}d = d$
- (d) $(d\mathcal{H})^2 = d\mathcal{H}$, $(\mathcal{H}d)^2 = \mathcal{H}d$.

Using (b), we observe that $\mathcal{H}d\omega \in \text{Ker } \mathcal{H}$. This and (14) can be used to define the *exact part* of ω as $\omega_e \equiv d\mathcal{H}\omega$ and the *anti-exact part* of ω as $\omega_a \equiv \omega - \omega_e = \mathcal{H}d\omega$.

Using (14), one can further show that $\mathcal{H}$ improves regularity. With no loss of generality, we prove this for forms having vanishing anti-exact part, *i.e.*, forms ω such that

$$\omega = \omega_e. \quad (15)$$

Note that for any given form ω , no cancelations can occur between ω_e and ω_a , therefore ω_e is always as smooth as ω , and furthermore $\mathcal{H}\omega_a = 0$. Thus we can

restrict our attention to forms satisfying (15). If ω is a one-form, then $\mathcal{H}\omega$ is a function and (15) implies that

$$\omega_i = \frac{\partial(\mathcal{H}\omega)}{\partial x^i} \forall i,$$

thus improving regularity. In the general case, $\mathcal{H}\omega \equiv (\mathcal{H}\omega)_I dx^I$, where I is a multi-index satisfying $|I| = k - 1$. In order to improve regularity in this case one would need to control $\delta(\mathcal{H}\omega)$, *i.e.*, all derivatives of type $\partial(\mathcal{H}\omega)_I / \partial x^i$ for $i \in I$. But $\delta(\mathcal{H}\omega) = 0$ from the Hodge Decomposition Theorem [25], as $\mathcal{H}\omega$ is anti-exact (using (b), above); so $\mathcal{H}$ is a smoothing operator on k -forms.

The important result for us is the following:

Recursive k -forms with coefficient Γ on a star-shaped region can be represented as follows [10]:

$$\omega = e^\eta [du + \mathcal{H}(\theta \wedge du)] , \quad (16)$$

where $\eta = \mathcal{H}\Gamma$, $\theta = \mathcal{H}d\Gamma$, and $u = \mathcal{H}(e^{-\eta}\omega)$. Condition (5) implies that θ satisfies

$$d\theta \wedge [du + \mathcal{H}(\theta \wedge du)] = 0.$$

For our purposes we can rewrite (16) as

$$\omega = e^\eta g(du), \quad (17)$$

where g is a smooth linear operator, or alternatively, as

$$\omega = e^\eta du + e^\eta h(du), \quad (18)$$

where h is also a smooth linear operator, the coefficients of which depend on Γ ; the latter variant yields better regularity. In fact, the form $h(du)$ is as smooth as u , provided Γ is smooth.

A particular case of the above occurs when the coefficient Γ is exact. In that case, the form ω satisfying (5) is said to be *gradient recursive*. For gradient recursive k -forms, (16) assumes the simpler form (10), in which g is the identity.

3 The linear case

Corresponding to the physical example of Sec. 1.2, which illustrates the non-linear form of eqs. (4), (5), we can illustrate the linear case by an even simpler physical example.

Condition (3) arises when a rigid body rotates in the xy -plane at constant angular velocity $\tilde{\omega}$. Taking the axis of rotation to lie at the origin of coordinates, we write the tangential velocity vector in the form

$$\mathbf{v} = v_1 \hat{i} + v_2 \hat{j},$$

where

$$v_1 = -\tilde{\omega}y, \quad v_2 = \tilde{\omega}x. \quad (19)$$

Then

$$\nabla \times \mathbf{v} = 2\tilde{\omega}\hat{k}, \quad (20)$$

so (3) is satisfied; see, *e.g.*, Exercise 4.4 of [24]. In the sequel we take $\tilde{\omega} \equiv 1$ for simplicity.

Equations (19) imply that

$$\nabla \cdot \mathbf{v} = 0,$$

so we express the 1-form ω dual to $\mathbf{v}$ as a solution of the *linear* Hodge-Frobenius equations

$$\begin{aligned} \delta\omega &= 0, \\ d\omega &= \Gamma \wedge \omega. \end{aligned} \quad (21)$$

Applying (10) and (13), we choose η to depend only on the distance r from the axis of rotation. Then

$$d\eta \wedge \omega = \eta'(r) \cdot r \, dxdy. \quad (22)$$

In addition, eq. (20) implies that

$$d\eta \wedge \omega = d\omega = 2 \, dxdy. \quad (23)$$

Equating the right-hand sides of eqs. (22) and (23), we conclude that $\eta(r) = 2 \log r$ and $\omega = r^2 du$ for $u(x, y) = \arctan(y/x)$. Then $|du| = r^{-1}$, so the singular structure of u in the xy -plane is analogous to that of the fundamental solution of Laplace's equation in $\mathbb{R}^3$. In particular, u is singular at the origin of the disc.

In this example, the Hodge-Frobenius equations themselves are only defined on the punctured disc, as

$$\Gamma \wedge \omega = (\Gamma_1 \omega_2 - \Gamma_2 \omega_1) \, dxdy = d\omega = 2 \, dxdy.$$

Using (19), we can write this condition as an equation for the inner product

$$\begin{pmatrix} \Gamma_1 \\ \Gamma_2 \end{pmatrix} \cdot \begin{pmatrix} x \\ y \end{pmatrix} = 2,$$

which cannot be satisfied at the origin.

Thus singular solutions arise naturally in both the linear and nonlinear Hodge-Frobenius equations.

Because eq. (16) requires that the domain be star-shaped, the conclusion that ω is representable as a product $f \, du$, where f is nonvanishing, does not follow and is in fact violated in our example, in which $f = r^2$.

We have presented a particularly simple model, in which the role of condition (5) is especially transparent. For more sophisticated completely integrable models of rigid-body rotation see, *e.g.*, [5] and references therein.

In the linear case we can accomplish easily what we cannot accomplish at all in the nonlinear case: an integrability condition sufficient to imply the smoothness of weak solutions.

Proposition 1. *Let ω be a weak solution of the linear Hodge-Frobenius equations (21) on Ω . If $|\Gamma|$ is bounded and $\omega \in L^p(\Omega)$ for $p > n$, then ω is continuous.*

Proof. The Friedrichs mollification ω_h of ω is a classical solution of (21); see, e.g., Sec. 7.2 of [13]. Thus

$$|d\omega_h|^p + |\delta\omega_h|^p + |\omega_h|^p \leq (|\Gamma|^p + 1) |\omega_h|^p.$$

Integrate and apply the L^p Gaffney-Gårding inequality (Lemma 4.7 of [16]) to obtain

$$\|\nabla\omega_h\|_p \leq C(\Gamma) \|\omega_h\|_p.$$

Because $\omega \in L^p(\Omega)$, we can allow the mollification parameter h to tend to zero. The proof is completed by the Sobolev Embedding Theorem.

4 Geometric analogies

4.1 Hypersurface-orthogonal vector fields

A unification of the two superficially different physical examples of Secs. 1 and 3 can be found in their underlying geometry – in particular, in the relation of each vector field to hypersurfaces created by the level sets of an associated scalar function u . We can think of u informally as the potential for the conservative field that would result from taking Γ to be zero in eq. (5). Whereas a conservative vector field is actually equal to ∇u , the vector fields in Secs. 1 and 3 merely point in the same direction as ∇u .

A nonvanishing vector field $\mathbf{v}$ is said to be *hypersurface-orthogonal* whenever there exists a foliation of hypersurfaces orthogonal to $\mathbf{v}$. The foliated hypersurfaces can be represented as level sets of a scalar function u . That is, one can write

$$\mathbf{v} = \lambda(\mathbf{x}) \nabla u$$

for a nonvanishing function λ . Conversely, a vector field which can be written in this way is clearly hypersurface-orthogonal. We conclude that a vector field $\mathbf{v}$ is hypersurface-orthogonal if and only if the 1-form ω dual to $\mathbf{v}$ satisfies

$$\omega = \lambda du ,$$

with nonvanishing λ (i.e., if and only if ω is completely integrable). Other equivalent conditions now follow from the Frobenius Theorem.

Hypersurface-orthogonal vector fields arise naturally in general relativity, particularly in connection with black-hole mechanics. Introducing a tensor field

$$\mathbf{B}_{\alpha\beta} \equiv \mathbf{v}_{\alpha;\beta},$$

where the semi-colon denotes covariant differentiation with respect to the space-time metric connection, the condition that $\mathbf{v}$ be hypersurface-orthogonal implies

that the antisymmetric part of $\mathbf{B}$, called the *rotation tensor*, vanishes ([29], Secs. 2.32, 2.33). For this reason, vector fields satisfying (5) are called *rotation-free* in general relativity, which is somewhat confusing in the context of the examples in Secs. 1 and 3.

4.2 Twisted Born-Infeld equations

A condition broadly analogous to (5) arises if ω is a Lie-algebra-valued 2-form satisfying the second Bianchi identity. In that case, replacing ω by F_A , where A is a Lie-algebra-valued 1-form, we have

$$dF_A = -[A, F_A], \quad (24)$$

where $[\cdot, \cdot]$ denotes the Lie bracket, an equation which resembles (5).

Precisely, let X be a vector bundle over a smooth, finite, oriented, n -dimensional Riemannian manifold M . Suppose that X has compact structure group $G \subset SO(m)$. Let $A \in \Gamma(M, ad X \otimes T^*M)$ be a connection 1-form on X having curvature 2-form

$$F_A = dA + \frac{1}{2}[A, A] = dA + A \wedge A,$$

where $[\cdot, \cdot]$ is the bracket of the Lie algebra $\mathfrak{S}$, the fiber of the adjoint bundle $ad X$. Sections of the automorphism bundle $Aut X$, called *gauge transformations*, act tensorially on F_A but affinely on A ; see, e.g., [22].

Consider energy functionals having the form (12), where $Q = |F_A|^2 = \langle F_A, F_A \rangle$ is an inner product on the fibers of the bundle $ad X \otimes \Lambda^2(T^*M)$. The inner product on $ad X$ is induced by the normalized trace inner product on $SO(m)$ and that on $\Lambda^2(T^*M)$, by the exterior product $*(F_A \wedge *F_A)$.

A nonabelian variational problem analogous to eqs. (5), (11) is described briefly in Sec. 5.1 of [28]. One is led to consider smooth variations taken in the infinitesimal deformation space of the connection and having the explicit form

$$\begin{aligned} var(E) &= \int_M \rho(Q) var(Q) dM = \int_M \rho(Q) \frac{d}{dt} \Big|_{t=0} |F_{A+t\psi}|^2 dM \\ &= \int_M \rho(Q) \frac{d}{dt} \Big|_{t=0} |F_A + tD_A\psi + t^2\psi \wedge \psi|^2 dM, \end{aligned} \quad (25)$$

where $D_A = d + [A, \cdot]$ is the exterior covariant derivative in the bundle. The Euler-Lagrange equations are

$$\delta(\rho(Q)F_A) = -*[A, *\rho(Q)F_A]. \quad (26)$$

In addition, we have the Bianchi identity (24).

Writing eq. (5) in components

$$d\omega^a = \Gamma_b^a \wedge \omega^b,$$

we observe that if $-\Gamma$ is interpreted as a connection 1-form, then (5) can be interpreted as the vanishing of an exterior covariant derivative, which is the content of eq. (24). Moreover, the well known algebraic requirement that Γ must satisfy

$$(d\Gamma_b^a - \Gamma_c^a \wedge \Gamma_b^c) \wedge \omega^b = 0$$

(*c.f.* eq. (4-2.3) of [10]) is the zero-curvature condition $[F_\Gamma, \omega] = 0$.

If G is abelian, then the Lie bracket vanishes and eqs. (26) reduce to the system

$$\delta \{ \rho [Q(F_A)] F_A \} = \delta \{ \rho [Q(dA)] dA \} = 0,$$

a nonlinear Hodge equation analogous to taking $\Gamma = 0$ in eqs. (4), (5).

Equations (24) reduce in the abelian case to the equations for the equality of mixed partial derivatives,

$$d^2 A = 0.$$

If $\rho \equiv 1$, then eqs. (26) are the Yang-Mills equations, describing quantum fields in the classical limit. These resemble a version of eq. (11) for 2-forms with $\rho \equiv 1$, with the Bianchi identity (24) playing the role of eq. (5).

Whereas the Yang-Mills equations do not have the nonlinear structure of (11) for non-constant ρ , those of the Born-Infeld model for electromagnetism are equivalent to (11) for differential forms of degree 2 with

$$\rho(Q) = (1 + |F_A|^2)^{-1/2}. \quad (27)$$

This model was introduced in [6] in order to produce a model of electromagnetism that does not diverge when the source is a point charge. Geometric aspects of the model are investigated in [11] and its analytic aspects, in [43]. A mathematical generalization of the Born-Infeld model to nonabelian variational equations was proposed in [26] and further studied in [38]; see also [15] for a related problem. The equations assume the form (24), (26) for an appropriate choice of $\rho(Q)$.

Geometrically, the nonabelian model puts a twist in the principal bundle corresponding to the configuration space of solutions. Thus eqs. (24), (26) are called *twisted nonlinear Hodge equations* [28]. A different approach to generalizing nonlinear Hodge theory to bundle-valued connections, which is based on the formulation of a natural class of boundary-value problems, is introduced in [23]; but interior estimates would be required in order to extend the theory of [23] to twisted forms of the equations considered in this paper. The derivation of such estimates is a goal of Sec. 5.2.

5 Analysis

We do not expect rotational fields of any kind to be very smooth. In particular, assumption (5) may produce caustics; see, for example, the discussion in Sec. 2.4 of [29]. However, it is reasonable to seek conditions under which the field

remains bounded at a singularity or under which the field equations remain uniformly elliptic.

Because these conditions will be derived for a large class of mass densities, the strength of the estimates obtained will depend on the integrability with respect to $\mathbf{x}$ of a given choice of mass density $\rho(Q(\mathbf{x}))$.

In the sequel we denote by C generic positive constants, the value of which may change from line to line. We follow an analogous convention for continually updated *small* positive constants ε . Repeated indices are to be summed from 1 to n .

5.1 Variational structure

The energy E of the field ω on Ω is defined by eq. (12), where $Q = Q(\omega)$ is defined as in (6) for $\omega \in \Lambda^1$ given by (10). Then $\forall \psi \in C_0^\infty(\Omega)$, the variations of E are computed as

$$\begin{aligned} \text{var}(E) &= \frac{d}{dt} E(u + t\psi)|_{t=0} = \frac{1}{2} \int_{\Omega} \rho(Q) \frac{d}{dt} Q(u + t\psi)|_{t=0} d\Omega \\ &= \frac{1}{2} \int_{\Omega} \rho(Q) \frac{d}{dt} [e^{2\eta} |d(u + t\psi)|^2]|_{t=0} d\Omega = \\ &= \frac{1}{2} \int_{\Omega} \rho(Q) e^{2\eta} \frac{d}{dt} (|du|^2 + 2\langle du, t d\psi \rangle + t^2 |d\psi|^2)|_{t=0} d\Omega = \\ &= \int_{\Omega} \rho(Q) e^{2\eta} \langle du, d\psi \rangle d\Omega = \int_{\Omega} \langle \rho(Q) e^{2\eta} du, d\psi \rangle d\Omega = \\ &= \int_{\Omega} d\langle \rho(Q) e^{2\eta} du, \psi \rangle d\Omega + \int_{\Omega} \langle \delta[\rho(Q) e^{\eta} \omega], \psi \rangle d\Omega = \int_{\Omega} \langle \delta[\rho(Q) e^{\eta} \omega], \psi \rangle d\Omega, \end{aligned}$$

as ψ has compact support in Ω . At a critical point, $\text{var}(E) = 0$, or

$$\delta[\rho(Q) e^{\eta} \omega] = 0, \quad (28)$$

so the variational formulation yields a “weighted” form of the continuity equation (4). The presence of this weight adds an inhomogeneous term to the unweighted variant. To see this, we write the local equation

$$-\partial_i [\rho(Q) e^{\eta} \omega_i] = -(\partial_i \eta) e^{\eta} \rho(Q) \omega_i - e^{\eta} \partial_i [\rho(Q) \omega_i] = 0.$$

This local form corresponds to the invariant representation

$$e^{\eta} \delta[\rho(Q) \omega] = e^{\eta} \rho(Q) \langle d\eta, \omega \rangle.$$

Thus we obtain eq. (11) for one-forms ω (i.e., for $k = 1$) or equivalently, since $\delta\alpha = - * d * \alpha$, $\forall \alpha \in \Lambda^1$, the equation

$$d * [\rho(Q) \omega] = -d\eta \wedge * [\rho(Q) \omega]. \quad (29)$$

5.1.1 Variational equations for k -forms

Recall that an exterior differential form ω of degree k which is recursive with coefficient Γ can be written in the form (18), where u is a $(k-1)$ -form (which depends on ω), and where the function η and the linear operator h depend only on Γ (Sec. 2.2).

We now compute the variation of the energy functional (12) among all forms $\omega + t\alpha$ satisfying

$$d(\omega + t\alpha) = \Gamma \wedge (\omega + t\alpha).$$

Such forms satisfy (18) with fixed η and h . Therefore the variations of E are computed as

$$\begin{aligned} \text{var}(E) &= \frac{d}{dt} E(\omega + t\alpha)|_{t=0} = \frac{1}{2} \int_{\Omega} \rho(Q) \frac{d}{dt} Q(\omega + t\alpha)|_{t=0} d\Omega \\ &= \int_{\Omega} \langle \rho(Q)\omega, \alpha \rangle d\Omega, \end{aligned}$$

where $\alpha = e^\eta (dv + h(dv))$ for a $(k-1)$ -form v which depends on α .

Thus

$$\begin{aligned} \text{var}(E) &= \frac{1}{2} \int_{\Omega} \langle e^\eta \rho(Q)\omega, dv + h(dv) \rangle d\Omega = \\ &= \int_{\Omega} \langle e^\eta \rho(Q)\omega, Gdv \rangle d\Omega = \int_{\Omega} \langle G^T e^\eta \rho(Q)\omega, dv \rangle d\Omega, \end{aligned}$$

where $G \equiv g_{ij}(\Gamma)$ is an $n \times n$ matrix and G^T is its transpose.

As the forms α (and thus v) are assumed to have compact support in Ω , setting $\text{var}(E) = 0$ is equivalent to imposing the condition

$$\delta [G^T e^\eta \rho(Q)\omega] = 0.$$

Notice that if ω is gradient recursive, then G is the identity matrix and we recover (28); or equivalently, since $\forall \alpha \in \Lambda^k$

$$\delta \alpha = (-1)^{nk+n+1} * d * \alpha, \quad (30)$$

the equation

$$d * [\rho(Q)\omega] = -d\eta \wedge * [\rho(Q)\omega], \quad (31)$$

i.e., eq. (11) for a gradient-recursive k -form ω with coefficient $d\eta$.

The variations employed in this section, applied directly to a 2-form, are necessarily different from the variations of (25), which are applied instead to the Lie-algebra valued connection 1-form A .

In the remainder of the paper we focus mainly on 1-forms. However, many of the results will extend easily to gradient-recursive k -forms. Moreover, the structure of expressions (17) and (18) suggest that, under appropriate technical hypotheses on the linear operators g or h , many of the results will extend also to recursive k -forms with coefficient Γ (not necessarily exact).

5.2 When are solutions bounded at a singularity?

Following [42], we find it convenient to introduce a function $H(Q)$ which is defined so that

$$H'(Q) = \frac{1}{2}\rho(Q) + Q\rho'(Q). \quad (32)$$

Then ellipticity is equivalent to the condition that H has positive derivative with respect to Q .

In Theorem 7 and Corollary 8 of [26], and in Theorem 6 and Corollary 7 of [27], L^p conditions are derived which imply the boundedness of solutions to eqs. (4), (5) on domains that include singular sets of given co-dimension. We call such theorems *partial removable singularities theorems*, as they imply that, although solutions may have jump discontinuities at the singularity, they cannot blow up there. Those results require the mass density ρ to satisfy the inequality

$$C(K + Q)^q \leq H'(Q) \leq C^{-1}(Q + K)^q \quad (33)$$

for constants $q > 0$ and $K \geq 0$. (This hypothesis is imposed in [27], following Sec. 1 of [42]; a somewhat stronger hypothesis is imposed in [26].)

There is obvious interest in deriving estimates for densities which may not satisfy (33). The focus of this section is to obtain and exploit such estimates for a broad class of densities, using the variational form of the equations. However, we retain the condition that ρ is positive, which is natural for applications.

We impose an additional condition that arises from technical considerations. If $\rho'(Q) > 0$, we require that

$$H'(Q) \leq C\rho(Q). \quad (34)$$

(This inequality is satisfied automatically if $\rho'(Q) \leq 0$.) If $\rho'(Q) < 0$ we require instead that

$$H'(Q) \geq C\rho(Q). \quad (35)$$

(This inequality is satisfied automatically if $\rho'(Q) \geq 0$.) Note that (35) implies (9) under our assumption $\rho(Q) > 0$. In the case $\rho'(Q) < 0$, eq. (11) is uniformly elliptic (i.e. (8) is satisfied) whenever (35) is satisfied and ρ is *noncavitating*: bounded below away from zero. In the case $\rho'(Q) > 0$, condition (8) is satisfied whenever $\rho(Q)$ is bounded above.

Densities which satisfy (33) satisfy the hypotheses (34), (35). However, there are many densities which satisfy (34), (35) but do not satisfy (33). Among the latter are densities for which the value of the exponent q on the left-hand side of inequality (33) differs from its value on the right-hand side, and certain densities for which the value of q in (33) is negative.

As a simple illustration, consider the class of densities

$$\rho(Q) = (K + Q)^q, \quad -1/2 < q < 0, \quad K > 0. \quad (36)$$

Such densities do not satisfy condition (33) and cavitate as Q tends to infinity. They arise, for example, in connection with models of pseudo-plastic

non-Newtonian fluids [2]. In this section we will obtain sufficient conditions for a bound on $Q\rho(Q)$ which is valid as Q tends to infinity. Under additional conditions, this bound extends to possibly singular solutions of the system (5), (11) with density given by (36). Note that for such densities, a bound on the product $Q\rho(Q)$ implies an asymptotic bound on the norm Q of the solution itself.

Because conditions (34) and (35) are used in a crucial way to establish both the subelliptic estimates and the ellipticity of the second-order operator, they appear to provide a mathematically natural generalization of condition (33).

Lemma 2. *Let condition (35) be satisfied. Then H can be chosen so that*

$$Q\rho(Q) \leq CH(Q). \quad (37)$$

Proof. If $\rho'(Q)$ is non-negative, then (37) is always satisfied with $C = 2$ if we choose $H(0) \geq 0$. (Note that assumption (35) is not needed in this case.) In order to see this, let

$$\Phi(Q) \equiv 2H(Q) - Q\rho(Q).$$

Then $\Phi(0) = H(0) \geq 0$, and

$$\Phi'(Q) = 2H'(Q) - \rho(Q) - Q\rho'(Q) = Q\rho'(Q) \geq 0.$$

Thus $\Phi(Q)$ remains nonnegative on the entire range of Q .

If $\rho'(Q) \leq 0$, we assume (35). Then in particular,

$$2H'(Q) \geq \varepsilon\rho(Q), \quad (38)$$

where we take ε to be so small that it lies in the interval $(0, 1)$. Inequality (38) can be written in the form

$$(1 - \varepsilon)\rho(Q) + 2Q\rho'(Q) \geq 0. \quad (39)$$

Define a constant c by the formula

$$c = \frac{1 + \varepsilon}{2\varepsilon}.$$

In terms of c , (39) can be written in the form

$$(c - 1)\rho(Q) + 2(c - 1/2)Q\rho'(Q) \geq 0.$$

We can convert this expression into the differential inequality

$$2cH'(s) \geq \frac{d}{ds} [s\rho(s)], \quad s \in [0, Q].$$

Integrate the inequality over s , using $H(0) = 0$. We obtain (37).

Lemma 3. *Let the 1-forms Γ and ω smoothly satisfy (5) and (11). Let $\rho > 0$ satisfy conditions (34) and (35). Then*

$$\Delta H + (-1)^{3n} \nabla \cdot \{ * [\omega \wedge * (\rho'(Q) dQ \wedge \omega)] \} + C (|\Gamma|^2 + |\nabla \Gamma|) H \geq 0. \quad (40)$$

Proof. We have (c.f. Sec. 1 of [42])

$$\begin{aligned} \langle \omega, \Delta [\rho(Q)\omega] \rangle &= \partial_i \langle \omega, \partial_i (\rho(Q)\omega) \rangle - \langle \partial_i \omega, \partial_i (\rho(Q)\omega) \rangle \\ &= \Delta H(Q) - [\rho(Q) \langle \partial_i \omega, \partial_i \omega \rangle + \rho'(Q) \langle \partial_i \omega, \omega \rangle \partial_i Q], \end{aligned} \quad (41)$$

where

$$\Delta H(Q) = \partial_i (\partial_i H(Q)) = \partial_i (H'(Q) \partial_i Q).$$

Writing $\partial_i Q = 2 \langle \partial_i \omega, \omega \rangle$, we rewrite (41) in the form

$$\langle \omega, \Delta [\rho(Q)\omega] \rangle = \Delta H(Q) - \rho(Q) |\nabla \omega|^2 - 2Q \rho'(Q) |d\omega|^2. \quad (42)$$

Applying eq. (42) to the operator identity $\Delta = -(\delta d + d\delta)$ and using (11), we write

$$\begin{aligned} 0 &= \langle \omega, \Delta [\rho(Q)\omega] \rangle + \langle \omega, \delta d (\rho(Q)\omega) \rangle + \langle \omega, d\delta (\rho(Q)\omega) \rangle \\ &= \Delta H(Q) - \gamma + \langle \omega, \delta (d\rho \wedge \omega) \rangle + \tau_1 + \tau_2, \end{aligned} \quad (43)$$

where

$$\begin{aligned} \gamma &= \rho(Q) |\nabla \omega|^2 + 2Q \rho'(Q) |d\omega|^2, \\ \tau_1 &= \langle \omega, \delta (\rho(Q) d\omega) \rangle, \end{aligned}$$

and

$$\tau_2 = \langle \omega, d [\rho(Q) \langle \Gamma, \omega \rangle] \rangle.$$

Define

$$L_\omega(H) \equiv \Delta H + \langle \omega, \delta (d\rho(Q) \wedge \omega) \rangle.$$

Then (43) can be written in the compact form

$$L_\omega H + \tau_1 + \tau_2 = \gamma. \quad (44)$$

We have

$$\begin{aligned} \tau_1 &= \langle \omega, \rho(Q) \delta (\Gamma \wedge \omega) - \langle d\rho(Q), d\omega \rangle \rangle \\ &\leq |\omega| [\rho(Q) |\delta (\Gamma \wedge \omega)| + |d\rho(Q)| |d\omega|] \leq \\ &|\omega| [\rho(Q) |\nabla \Gamma| |\omega| + \rho(Q) |\Gamma| |\nabla \omega| + |\rho'(Q) dQ| |d\omega|]. \end{aligned}$$

Applying (5) to the last term on the right-hand side and using

$$|dQ| = 2|\omega| |d\omega|,$$

we obtain

$$\begin{aligned}\tau_1 \leq & Q\rho(Q)|\nabla\Gamma| + \rho(Q)|\Gamma||\omega||\nabla\omega| + 2Q|\rho'(Q)||\Gamma||\omega||d|\omega| \leq \\ & Q\rho(Q)|\nabla\Gamma| + \rho(Q) \left[\frac{Q|\Gamma|^2}{2\varepsilon} + \frac{\varepsilon}{2}|\nabla\omega|^2 \right] \\ & + 2Q|\rho'(Q)| \left(\frac{\varepsilon}{2}|d|\omega||^2 + \frac{|\Gamma|^2 Q}{2\varepsilon} \right).\end{aligned}\quad (45)$$

Estimating τ_2 yields the same terms:

$$\begin{aligned}\tau_2 \leq & Q\rho(Q)|\nabla\Gamma| + |\omega||\Gamma|[\rho(Q)|\nabla\omega| + |\omega||\nabla\rho(Q)|] = \\ & Q\rho(Q)|\nabla\Gamma| + |\omega||\Gamma|[\rho(Q)|\nabla\omega| + 2Q|\Gamma||\rho'(Q)||\omega||d|\omega|],\end{aligned}\quad (46)$$

which is bounded by the right-hand side of (45). That is,

$$\begin{aligned}\tau_1 + \tau_2 \leq & 2Q\rho(Q)|\nabla\Gamma| + \frac{Q}{\varepsilon}[\rho(Q) + 2Q|\rho'(Q)|]|\Gamma|^2 + \\ & \varepsilon\rho(Q)|\nabla\omega|^2 + 2Q\varepsilon|\rho'(Q)||d|\omega||^2.\end{aligned}\quad (47)$$

Applying inequality (47) to eq. (44), we obtain, in the case $\rho'(Q) > 0$, the estimate

$$\begin{aligned}L_\omega(H) + 2 \left[Q\rho(Q)|\nabla\Gamma| + \frac{QH'(Q)}{\varepsilon}|\Gamma|^2 \right] \geq \\ (1 - \varepsilon) [\rho(Q)|\nabla\omega|^2 + 2Q\rho'(Q)|d|\omega||^2] \geq (1 - \varepsilon) H'(Q)|d|\omega||^2,\end{aligned}\quad (48)$$

the inequality on the right following from Kato's inequality. We apply Lemma 2 and (34) to terms on the extreme left-hand side of inequality (48):

$$\begin{aligned}2 \left[Q\rho(Q)|\nabla\Gamma| + \frac{QH'(Q)}{\varepsilon}|\Gamma|^2 \right] \leq \\ C [H(Q)|\nabla\Gamma| + Q\rho(Q)|\Gamma|^2] \leq \\ C (|\nabla\Gamma| + |\Gamma|^2) H(Q).\end{aligned}$$

We now have, for the case $\rho'(Q) > 0$, the estimate

$$L_\omega(H) + C (|\nabla\Gamma| + |\Gamma|^2) H \geq (1 - \varepsilon) H'(Q)|d|\omega||^2 \geq 0. \quad (49)$$

In the case $\rho'(Q) < 0$, we also apply (47) to (44); but here we obtain

$$\begin{aligned}L_\omega(H) + 2 \left[Q\rho(Q)|\nabla\Gamma| + \frac{QH'(Q)}{\varepsilon}|\Gamma|^2 \right] \geq \\ (1 - \varepsilon) [\rho(Q)|\nabla\omega|^2 + 2Q\rho'(Q)|d|\omega||^2] + 4\varepsilon Q\rho'(Q)|d|\omega||^2.\end{aligned}\quad (50)$$

In the case $\rho'(Q) < 0$, condition (35) yields

$$-Q\rho'(Q) < \frac{1}{2}\rho(Q) \leq CH'(Q),$$

thus

$$Q|\rho'(Q)| \leq CH'(Q). \quad (51)$$

(Note that (51) is automatic for $\rho'(Q) \geq 0$.) Applying Kato's inequality and (51) to (50) yields

$$L_\omega(H) + 2 \left[Q\rho(Q)|\nabla\Gamma| + \frac{QH'(Q)}{\varepsilon}|\Gamma|^2 \right] \geq \\ (1 - \varepsilon - 4\varepsilon C) H'(Q)|d\omega|^2 \geq 0.$$

Applying (34) and (37) to the left-hand side of this inequality, we obtain (49) (for updated ε) for the case $\rho'(Q) < 0$ as well.

It remains only show that the operator $L_\omega(H)$ can be put into divergence form, at the cost of absorbing another lower-order term. The second term of $L_\omega(H)$ can be written in the form

$$\langle \omega, \delta(d\rho(Q) \wedge \omega) \rangle = *[\omega \wedge * \delta(d\rho(Q) \wedge \omega)] = \\ *d[\omega \wedge *(d\rho(Q) \wedge \omega)] - *[d\omega \wedge *(d\rho(Q) \wedge \omega)],$$

where

$$-*[d\omega \wedge *(d\rho(Q) \wedge \omega)] = -*[\Gamma \wedge \omega \wedge *(d\rho(Q) \wedge \omega)] \geq \\ -2Q|\Gamma||\rho'(Q)||\omega||d\omega|,$$

which is estimated in the same way as the last term in the sum on the extreme right-hand side of (46).

Taking into account (30) we obtain, for any k -form α ,

$$*d\alpha = (-1)^{k(n-k)} *d(**)\alpha = \\ (-1)^{k(n-k)} (*d*)\alpha = (-1)^{2kn+n+1-k^2} \delta * \alpha. \quad (52)$$

Taking $k = 1$ and

$$\alpha = *[\omega \wedge *(\rho'(Q)dQ \wedge \omega)]$$

in eq. (52), we can express the operator δ in that equation as a divergence. This allows us to write (49) (again updating C and ε) in the form

$$\Delta H + (-1)^{3n} \nabla \cdot \{ *[\omega \wedge *(\rho'(Q)dQ \wedge \omega)] \} + C(|\Gamma|^2 + |\nabla\Gamma|) H \geq \\ (1 - \varepsilon) H'(Q)|d\omega|^2 \geq 0.$$

This completes the proof of Lemma 3.

Lemma 4. *Under the hypotheses of Lemma 3, the operator*

$$\mathcal{L}_\omega(H) \equiv \Delta H + (-1)^{3n} \nabla \cdot \{ *[\omega \wedge *(\rho'(Q)dQ \wedge \omega)] \}$$

is a uniformly elliptic operator on H .

Proof. We argue as in Sec. 1 of [42] and Sec. 4 of [27], but without using hypothesis (33).

Define a map $\beta_\omega : \Lambda^0 \rightarrow \Lambda^{k+1}$ by the explicit formula

$$\beta_\omega : \mu \rightarrow d\mu \wedge \omega, \quad (53)$$

for $\mu \in \Lambda^0$ and $\omega \in \Lambda^k$. Then we can write the variational form of the Hodge-Frobenius equations (5), (10), (11) in the alternate form

$$d\sigma_i = -\beta_{\sigma_i}(\eta), \quad \sigma = 1, 2, \quad (54)$$

where $\sigma_1 = *\rho(Q)\omega$ and $\sigma_2 = -\omega$. The “irrotational” case $d\omega = 0$ can be recovered as the special case of (54) in which

$$\beta_\omega(\eta) = d(\eta\omega),$$

in which case eq. (53) implies that

$$d\eta \wedge \omega = d\eta \wedge \omega + \eta d\omega$$

(c.f. Sec. 4 of [27] and Sec. 1 of [42]).

Moreover, writing

$$\beta_\omega(g) = dg \wedge \omega$$

for some 0-form g , we compute for arbitrary compactly supported $\mu \in \Lambda^{k+1}$,

$$\langle \mu, dg \wedge \omega \rangle = *(dg \wedge (**)(\omega \wedge *\mu)) = \langle dg, *(\omega \wedge *\mu) \rangle = \langle g, \delta *(\omega \wedge *\mu) \rangle.$$

So the map $\beta_\omega^* : \Lambda^{k+1} \rightarrow \Lambda^0$ defined by the explicit formula

$$\beta_\omega^*(\mu) = \delta *(\omega \wedge *\mu)$$

is the formal adjoint of β_ω .

In terms of the maps β_ω and β_ω^* , we can write

$$\begin{aligned} (-1)^{n+1} \nabla \cdot \{ *[\omega \wedge *(\rho'(Q)dQ \wedge \omega)] \} &= \beta_\omega^* \beta_\omega[\rho] \\ &= \beta_\omega^*[\mu_\omega(H)] \end{aligned}$$

for μ_ω satisfying

$$\mu_\omega(H) = \frac{\rho'(Q)}{H'(Q)} dH \wedge \omega.$$

Using β_ω^* , we write the inequality of Lemma 3 in the form

$$\mathcal{L}_\omega(H) + \text{lower-order terms} \geq 0,$$

where

$$\mathcal{L}_\omega(H) = \Delta H - \beta_\omega^*[\mu_\omega(H)].$$

Writing

$$\mathcal{L}_\omega(H) = \partial_k (\alpha^{jk} \partial_j) H,$$

we find that if $\rho'(Q) < 0$, then (51) implies that

$$1 \leq \alpha_{kj} + \frac{Q|\rho'(Q)|}{H'(Q)} \leq \alpha_{kj} + C.$$

If $\rho'(Q) \geq 0$, then we write

$$\nabla \cdot \left\{ \left[1 - \frac{Q\rho'(Q)}{H'(Q)} \right] \nabla H \right\} = \nabla \cdot \left\{ \left[\frac{\rho(Q)}{2H'(Q)} \right] \nabla H \right\};$$

condition (34) implies that there is a positive constant c such that

$$\frac{c}{2} \leq \frac{\rho(Q)}{2H'(Q)} \leq 1;$$

c is the reciprocal of the constant C in (34). This completes the proof of Lemma 4.

These three lemmas easily yield:

Theorem 5. *Under the hypotheses of Lemma 3, the product $Q\rho(Q)$ is locally bounded above by the L^2 -norm of H .*

The proof of Theorem 5 is given in Sec. 7.1.

Of course the integrability of H depends on ρ . But for any given ρ in $C^1(Q)$, H can be computed explicitly by integrating (32). Even if ρ cavitates, Theorem 5 yields asymptotic information about the fastest rate at which Q can blow up. Nevertheless, Theorem 5 is ultimately not very useful, due to the hypothesis that the solutions are smooth. It would become more useful if it could be applied to singular solutions. We will find that the partial removable singularities theorems proven in [27] for singular sets of prescribed codimension extend to our conditions on ρ under slightly different hypotheses.

Initially, we treat the special case of an isolated point singularity, for which the proof is somewhat simpler than the proof for higher-order singularities and the range of applicable dimensions somewhat larger.

Theorem 6. *Let the hypotheses of Lemma 3 be satisfied on $\Omega \setminus \{p\}$, where p is a point of $\mathbb{R}^n$ and $n > 2$. If $H \in L^{2n/(n-2)}(\Omega)$ and if the function*

$$f \equiv |\nabla \Gamma| + |\Gamma|^2 \tag{55}$$

is sufficiently small in $L^{n/2}(\Omega)$, then H is an $H^{1,2}$ -weak solution in a neighborhood of the singularity.

The proof of Theorem 6 is given in Sec. 7.2.

Corollary 7. *Let the hypotheses of Theorem 6 be satisfied and, in addition, let the function f given by (55) satisfy the growth condition*

$$\int_{B_r(\mathbf{x}_0) \cap \Omega} |f|^{n/2} d\Omega \leq Cr^\kappa \quad (56)$$

for some $\kappa > 0$, where $B_r(\mathbf{x}_0)$ is an n -disc of radius r , centered at $\mathbf{x}_0$. Then the conclusion of Theorem 5 remains valid.

Proof. Apply Theorem 5.3.1 of [25] to the conclusion of Theorem 6, following the proof of Theorem 5.

Note that the singularity in Corollary 7 is in the solution, rather than in the underlying metric (*c.f.* [39], in which metric point singularities are considered in the case $\Gamma \equiv 0$). Corollary 7 extends to higher-order singularities in spaces of sufficiently high dimension; but the proof requires more delicate test functions.

Theorem 8. *Let the pair ω, Γ smoothly satisfy eqs. (5), (11), with ρ satisfying the hypotheses of Lemma 3, on $\Omega \setminus \Sigma$. Here Σ is a compact singular set, of dimension $0 \leq m < n - 4$, completely contained in a sufficiently small n -disc D which is itself completely contained in the interior of Ω . Let $H(Q)$ lie in $L^{2\beta\gamma_1}(D) \cap L^{2\gamma_2}(D)$, where $\beta = (n - m - \varepsilon) / (n - m - 2 - \varepsilon)$ for $1/2 < \gamma_1 < \gamma_2$. If the function f given by (55) satisfies the growth condition (56), then the conclusion of Theorem 5 remains valid.*

The proof of Theorem 8 is given in Sec. 7.3.

The conditions imposed on ρ in this section also lead to extensions of known results for the conventional case $\Gamma \equiv 0$. In particular, we consider equations which, expressed in components, have the weak form

$$\int_{\Omega} [\rho(Q)u_{x_k}]_{x_i} \varphi_{x_k} d\Omega = 0, \quad (57)$$

where $\varphi \in C_0^\infty(\Omega)$ is arbitrary and $Q = |du|^2$. Such equations have been intensively studied in cases for which $\rho(|du|^2)$ grows as a power of $|du|$; *c.f.* eq. (3.10) of [8]. Equation (57) can be interpreted as a weak derivative with respect to x_i of the system (4), (5) with $\Gamma \equiv 0$.

Define

$$\mathfrak{H}(Q) \equiv Q\rho^2(Q).$$

The system (4), (5) is elliptic precisely when $\mathfrak{H}'(Q) > 0$. The following result extends Theorem 1 of [27], which requires the derivative of ρ with respect to Q to be nonpositive. If ρ is specified to be nonincreasing in Q , then the integrability of $\mathfrak{H}(Q)$ can be shown to follow from finite energy. In the general case, we impose this integrability as an independent hypothesis on the weak solution.

Theorem 9. *Let the scalar function $u(\mathbf{x})$ satisfy equation (57) with ρ bounded, positive, and noncavitating, and with $\mathfrak{H}(Q)$ integrable. Assume conditions (9) and (34). Then for every n -disc D_R of radius R completely contained in Ω there is a positive number $\delta > 0$ such that*

$$\sup_{Q \in D_{(1-\delta)R}} \mathfrak{H}(Q) \leq CR^{-n} \int_{D_R} \mathfrak{H}(Q) * 1,$$

where C depends on ρ and δ but not on Q or R .

Proof. Define Q_* to be the (possibly infinite) value of Q for which

$$\sup_{Q \in \Omega} \rho(Q) = \rho(Q_*).$$

Then

$$\frac{\mathfrak{H}'(Q)}{\rho^2(Q_*)} = \frac{2\rho(Q)H'(Q)}{\rho^2(Q_*)} \leq C \left(\frac{\rho(Q)}{\rho(Q_*)} \right)^2 \leq C, \quad (58)$$

where C is the constant of (34).

As in [27], we initially estimate smooth solutions, and subsequently extend the result to weak solutions, recovering the derivatives as limits of difference quotients.

We choose test functions φ^i having the form

$$\varphi^i(\mathbf{x}) = u_{x_i} \tilde{\mathfrak{H}}^{\alpha/2} \zeta^2,$$

where $\alpha > 0$, $\zeta(\mathbf{x}) \in C_0^\infty(D_R)$, and

$$\tilde{\mathfrak{H}} \equiv \mathfrak{H}(Q) + \varepsilon \quad (59)$$

for a small positive parameter ε (c.f. [8], Sec. 3). Then

$$\begin{aligned} & [\rho(Q)u_{x_k}]_{x_i} \varphi_{x_k}^i = \\ & 2H'(Q)(u_{x_i x_k})^2 \tilde{\mathfrak{H}}^{\alpha/2} \zeta^2 + \alpha \rho(Q) [H'(Q)]^2 |\nabla Q|^2 \tilde{\mathfrak{H}}^{(\alpha-2)/2} \zeta^2 \\ & + 2H'(Q)Q_{x_k} \tilde{\mathfrak{H}}^{\alpha/2} \zeta \zeta_{x_k} \equiv i_1 + i_2 + i_3. \end{aligned}$$

Here

$$\begin{aligned} i_1 & \geq 2H'(Q)u_{x_i x_k} u_{x_i x_k} Q \rho^2(Q) \tilde{\mathfrak{H}}^{(\alpha-2)/2} \zeta^2 = \\ & \frac{1}{2} \rho^2(Q) H'(Q) |\nabla Q|^2 \tilde{\mathfrak{H}}^{(\alpha-2)/2} \zeta^2 \geq C |\nabla (\tilde{\mathfrak{H}}^{(\alpha+2)/4})|^2 \zeta^2, \end{aligned}$$

where the last inequality follows from (58) and the constant C depends on α , $\rho^{-1}(Q_*)$, the constant of (58), and the lower bound of $\rho(Q)$.

Similarly,

$$i_2 \geq \frac{\rho(Q)}{\rho(Q_*)} i_2 \geq C |\nabla (\tilde{\mathfrak{H}}^{(\alpha+2)/4})|^2 \zeta^2$$

and

$$i_3 = 2H'(Q)Q_{x_j}\tilde{\mathfrak{H}}^{\alpha/2}\zeta(\mathbf{x})\zeta_{x_j}.$$

The latter quantity can be estimated by applying, as in the proof of Lemma 3, the elementary algebraic inequality (*Young's inequality*)

$$ab \geq -\left(\frac{\tilde{\varepsilon}}{2}a^2 + \frac{1}{2\tilde{\varepsilon}}b^2\right),$$

now taking

$$a = 2H'(Q)\tilde{\mathfrak{H}}^{(\alpha-2)/4}\zeta Q_{x_j}, \quad b = \tilde{\mathfrak{H}}^{(\alpha+2)/4}\zeta_{x_j},$$

and $\tilde{\varepsilon} = \tilde{\delta}\rho^2(Q)$ for small $\tilde{\delta} > 0$.

Putting these estimates together, we obtain

$$\int_{\Omega} |\nabla \left(\tilde{\mathfrak{H}}^{(\alpha+2)/4} \right)|^2 \zeta^2 d\Omega \leq C \int_{\Omega} \tilde{\mathfrak{H}}^{(\alpha+2)/2} |\nabla \zeta|^2 d\Omega.$$

The proof for the smooth case is completed by applying the Moser iteration as in expressions (9.5.8)-(9.5.12) of [18] and subsequently letting the parameter ε in (59) tend to zero. The proof is extended to the general case by applying the difference-quotient method as in eq. (13) of [27]; see also Lemma 2 of [42].

6 Hodge-Bäcklund transformations

Different choices of ρ may sometimes be related by a special kind of Bäcklund transformation which is based on properties of the Hodge involution. We call these transformations *Hodge-Bäcklund*. Although this term does not seem to have been used up to now, such generalized Bäcklund transformations have a long history in diverse fields of mathematical physics. Our aim in this section is to unify these various transformations, place them in an invariant context, and extend them to the completely integrable case.

Historically, the term *Bäcklund transformation* has been defined in many ways; see, *e.g.*, [31] for the classical theory. In the sequel we will use it in the general sense of a function that maps a solution a of a differential equation A into a solution b of a differential equation B and vice-versa, where B may equal A but b will not equal a .

6.1 Transformation of the Chaplygin mass density

The mass density for the adiabatic and isentropic subsonic flow of an ideal fluid has the form

$$\rho(Q) = \left(1 - \frac{\gamma-1}{2}Q\right)^{1/(\gamma-1)}, \quad (60)$$

for $Q \in [0, 2/(\gamma+1))$, where γ is the *adiabatic constant*: the ratio of specific heats for the gas. The adiabatic constant for air is 1.4. Choosing γ to be 2 we obtain, by an independent physical argument originally introduced for one

space dimension in [30], the mass density for shallow hydrodynamic flow in the tranquil regime (*c.f.* eq. (10.12.5) of [40]). If we choose γ to be -1 (a physically impossible choice), we obtain the density of the minimal surface equation [17], [37]

$$\rho(Q) = \frac{1}{\sqrt{1+Q}} \quad (61)$$

Flow governed by this density is called *Chaplygin flow*. Despite the fact that the numbers -1 and 1.4 are not particularly close, this choice of mass density has many attractive properties as an approximation for (60); see, *e.g.*, [9] and Ch. 5 of [4]. These properties are, in general, retained in the case of completely integrable flow described in Sec. 1.2.

If $\Gamma \equiv 0$, eqs. (4) and (5) with $\rho(Q)$ given by (61) describe, for $k = 1$, nonparametric minimal surfaces embedded in Euclidean space. If $k = 2$ they describe electromagnetic fields in the Born-Infeld model, as in (27).

More generally, we have the following result, which extends an argument introduced for the case $\Gamma \equiv 0$ by Yang in [43]; see also [1] and Theorem 2.1 of [38].

Theorem 10. *Let the 1-form ω satisfy eqs. (5) and (11), with ρ satisfying (61). Then there exists an $(n-1)$ -form ξ with $|\xi| < 1$, satisfying equations analogous to (5) and (11), but with $\Gamma \equiv d\eta$ replaced by $\hat{\Gamma} \equiv d\hat{\eta} = -d\eta$ and $\rho(Q)$ replaced by*

$$\hat{\rho}(|\xi|^2) \equiv \frac{1}{\sqrt{1-|\xi|^2}}. \quad (62)$$

Proof. Equation (11) can be interpreted as the assertion that the $(n-1)$ -form

$$\xi = *[\rho(Q)\omega] = * \left[\frac{\omega}{\sqrt{1+|\omega|^2}} \right] \quad (63)$$

satisfies

$$d\xi = d\hat{\eta} \wedge \xi, \quad (64)$$

that is, equation (5) with η replaced by $\hat{\eta} \equiv -\eta$. As a consequence, we conclude that this $(n-1)$ -form is also gradient recursive and, on domains with trivial de Rham cohomology, there exists an $(n-2)$ -form σ such that $\xi = e^{\hat{\eta}} d\sigma$ (*c.f.* Sec. 2.2, final paragraph).

Because the Hodge involution is an isometry,

$$|\xi|^2 = \frac{|\omega|^2}{1+|\omega|^2}$$

or, equivalently,

$$1 - |\xi|^2 = \frac{1}{1+|\omega|^2}. \quad (65)$$

Note that eq. (65) implies $|\xi|^2 < 1$, as well as

$$\rho(|\omega|^2)\hat{\rho}(|\xi|^2) = \frac{1}{\sqrt{1+|\omega|^2}} \frac{1}{\sqrt{1-|\xi|^2}} = 1.$$

This, together with (63), yields directly

$$*\hat{\rho}(|\xi|^2)\xi = *^2\rho(|\omega|^2)\hat{\rho}(|\xi|^2)\omega = (-1)^{n-1}\omega.$$

Hence,

$$d*(\hat{\rho}(|\xi|^2)\xi) = (-1)^{n-1}d\omega = (-1)^{n-1}d\eta\wedge\omega = d\eta\wedge*(\hat{\rho}(|\xi|^2)\xi) = -d\hat{\eta}\wedge*(\hat{\rho}(|\xi|^2)\xi),$$

which is equivalent to equation (11) for the gradient recursive $(n-1)$ -form ξ with coefficient $d\hat{\eta}$, where ρ has been replaced by $\hat{\rho}$ (see also (31)). This completes the proof of Theorem 10.

The above argument carries over to any pairing of functions $\rho(|\omega|^2)$, $\hat{\rho}(|\xi|^2)$, as long as their product is 1. Moreover, it carries over essentially unchanged to gradient-recursive k -forms ω , as these would automatically yield gradient-recursive $(n-k)$ -forms ξ .

The same argument, with small modifications, extends Theorem 10 to general (non-gradient-recursive) k -forms, in which we write equations (5) and (11) in terms of Γ and $\hat{\Gamma}$ rather than in terms of η and $\hat{\eta}$.

In addition to the original, “irrotational” version of Theorem 10 introduced in [43], other aspects of the duality of mass densities for nonlinear Hodge equations are presented in [36], and in Sec. 4.2 of [28]. (The irrotational case of the above argument is reviewed in Sec. 4.1 of [28]; note the recurring misprint in the two paragraphs following eq. (35) of that reference: $d\omega$ should be du .) Densities of the form (62) arise in the study of maximal space-like hypersurfaces [7] and, in a completely different way, harmonic diffeomorphisms [41].

6.2 Transformation of the complex eikonal equation (*after Magnanini and Talenti*)

As an example of the diverse fields in which these transformations arise, and of the simplifying and unifying role of the Hodge involution, we describe an example from complex optics. The description of the local Bäcklund transformations follows the analysis of Magnanini and Talenti [19], in which these transformations were introduced; see also [20] and [21]. We then reproduce the argument of [19] in a simpler, invariant context using the Hodge operator.

The *eikonal equation* in $\mathbb{R}^2$ can be written in the form

$$\psi_x^2 + \psi_y^2 + \nu^2 = 0, \tag{66}$$

where $\nu(x, y)$ is a given real-valued function. If we write the solution $\psi(x, y)$ as a complex function having the form

$$\psi(x, y) = u(x, y) + iv(x, y)$$

for real-valued functions u and v , then (66) is equivalent to the first-order system

$$u_x^2 + u_y^2 - v_x^2 - v_y^2 + \nu^2 = 0, \quad (67)$$

$$u_x v_x + u_y v_y = 0. \quad (68)$$

The function ν corresponds physically to the refractive index of the medium through which the wavefront represented by the function ψ propagates. If

$$u_x^2 + u_y^2 > 0, \quad (69)$$

then v_x and v_y cannot both vanish, by eq. (67). In that case, eq. (68) can be expressed, in the language of proportions, by the assertion that either

$$\begin{bmatrix} v_x \\ v_y \end{bmatrix} : \sqrt{v_x^2 + v_y^2} = \pm \begin{bmatrix} -u_y \\ u_x \end{bmatrix} : \sqrt{u_x^2 + u_y^2}, \quad (70)$$

or

$$u_x = u_y = 0. \quad (71)$$

Treating these relations as a coupled system of scalar equations, we have, under the same hypothesis,

$$\frac{v_x}{\sqrt{v_x^2 + v_y^2}} = \pm \frac{-u_y}{\sqrt{u_x^2 + u_y^2}}$$

and, using eq. (67),

$$v_x = \pm (-u_y) \sqrt{\frac{\nu^2}{u_x^2 + u_y^2} + 1}. \quad (72)$$

Similarly, the scalar equation

$$\frac{v_y}{\sqrt{v_x^2 + v_y^2}} = \pm \frac{u_x}{\sqrt{u_x^2 + u_y^2}},$$

which also follows from (70), implies by analogous operations the equation

$$v_y = \pm u_x \sqrt{\frac{\nu^2}{u_x^2 + u_y^2} + 1}. \quad (73)$$

We can write the coupled system (72), (73) as a vector equation of the form

$$\begin{bmatrix} v_x \\ v_y \end{bmatrix} = \pm \sqrt{1 + \frac{\nu^2}{u_x^2 + u_y^2}} \begin{bmatrix} -u_y \\ u_x \end{bmatrix}, \quad (74)$$

or as a single (exact) equation for 1-forms,

$$dv = \pm \sqrt{1 + \frac{\nu^2}{u_x^2 + u_y^2}} (-u_y dx + u_x dy). \quad (75)$$

This implies the local existence of a solution to the divergence-form equation

$$\frac{\partial}{\partial x} \left(\sqrt{1 + \frac{\nu^2}{u_x^2 + u_y^2}} u_x \right) + \frac{\partial}{\partial y} \left(\sqrt{1 + \frac{\nu^2}{u_x^2 + u_y^2}} u_y \right) = 0, \quad (76)$$

whenever condition (69) is satisfied.

The equations (67), (68), and (70) can also be solved for u_x and u_y , in addition to being solvable for v_x and v_y as in (72) and (73). Under the hypothesis that either

$$v_x^2 + v_y^2 = \nu^2 \quad (77)$$

or

$$v_x^2 + v_y^2 > \nu^2, \quad (78)$$

one obtains, by completely analogous arguments to those applied to v_x and v_y , the vector equation

$$\begin{bmatrix} u_x \\ u_y \end{bmatrix} = \mp \sqrt{1 - \frac{\nu^2}{v_x^2 + v_y^2}} \begin{bmatrix} -v_y \\ v_x \end{bmatrix} \quad (79)$$

and the equation

$$du = \mp \sqrt{1 - \frac{\nu^2}{v_x^2 + v_y^2}} (-v_y dx + v_x dy), \quad (80)$$

for 1-forms. Note that eq. (80) is exact if either (77) holds, or if (78) holds and the divergence-form equation

$$\frac{\partial}{\partial x} \left(\sqrt{1 - \frac{\nu^2}{v_x^2 + v_y^2}} v_x \right) + \frac{\partial}{\partial y} \left(\sqrt{1 - \frac{\nu^2}{v_x^2 + v_y^2}} v_y \right) = 0 \quad (81)$$

is satisfied.

6.2.1 Hodge-Bäcklund interpretation

Equations (76) and (81) define a Bäcklund transformation $u \rightarrow v$ and its inverse $v \rightarrow u$ (*c.f.* [21]). These equations can be written in the form of nonlinear Hodge equations

$$d * (\rho(Q)\omega) = 0, \quad (82)$$

$$d\omega = 0 \quad (83)$$

(that is, as eqs. (4), (5) for $\Gamma \equiv 0$). Either

$$\rho(|\omega|^2) = \sqrt{1 + \frac{\nu^2}{|\omega|^2}} \quad (84)$$

corresponding to eq. (76), or

$$\hat{\rho}(|\xi|^2) = \sqrt{1 - \frac{\nu^2}{|\xi|^2}} \quad (85)$$

corresponding to eq. (81). In either case we assume that Q does not vanish, by analogy with the equations (69), (71), (77), and (78).

Take ρ as in (84). If the domain is simply connected, then eq. (83) implies, analogously to Sec. 5.1, that there is a 0-form τ such that

$$\omega = du,$$

and a 0-form v such that

$$dv = \pm * \left(\sqrt{1 + \frac{\nu^2}{|\omega|^2}} \omega \right) = \pm * \left(\sqrt{|du|^2 + \nu^2} \frac{\omega}{|\omega|} \right). \quad (86)$$

Because the Hodge operator is an isometry – which is illustrated locally by (70), we have

$$|dv|^2 = |du|^2 + \nu^2.$$

Thus the Hodge-Bäcklund transformation (86) yields an invariant form of (67). Unlike classical Bäcklund transformations of the eikonal equation, in this case the Cauchy-Riemann equations are not satisfied. Rather,

$$u_x = \mp \rho(Q) v_y$$

and

$$u_y = \pm \rho(Q) v_x,$$

which is sufficient for the orthogonality condition (68).

Now take $\hat{\rho}$ as in (85). Arguing as before, we conclude that there is a 0-form $\tilde{u}$ such that $\omega = d\tilde{u}$, and a 0-form $\tilde{v}$ such that

$$d\tilde{v} = \pm * \left(\sqrt{Q - \nu^2} \frac{\omega}{|\omega|} \right).$$

We obtain

$$|d\tilde{v}|^2 = |d\tilde{u}|^2 - \nu^2.$$

Letting $\tilde{u} = \pm iu$ and $\tilde{v} = \pm iv$, we obtain a mapping taking solutions to eqs. (5), (11) with $\Gamma \equiv 0$ and $\hat{\rho}$ satisfying (85) into solutions of that system with $\Gamma \equiv 0$ and ρ satisfying (84).

These arguments extend immediately to the Hodge-Frobenius case – for example, by replacing (86) with the expansion

$$e^{\hat{\eta}} dv = \pm * \left(\sqrt{e^{2\eta} |du|^2 + \nu^2} \frac{\omega}{|\omega|} \right),$$

and squaring both sides. They also extend in a straightforward way to gradient-recursive k -forms and, with some modifications, to general k -forms (see the remarks following Theorem 10).

Motivated by these examples, we offer a general definition of the *Hodge-Bäcklund transformation*. It is a map taking a solution a of a nonlinear Hodge-Frobenius equation having mass density ρ_A into a solution b of a nonlinear Hodge-Frobenius equation having mass density ρ_B and vice-versa, where B may equal A but b will not equal a .

7 Appendix: Methods from elliptic theory

In this appendix we collect the proofs of Theorem 5, 6, and 8, which follow directly from the association of solutions to a uniformly subelliptic operator via Lemmas 2-4.

7.1 Proof of Theorem 5

We require the following well known extension of de Giorgi-Nash-Moser theory:

Theorem A.1 (Morrey [25], Theorem 5.3.1). *Let $n > 2$. Let $U \in H^{1,2}(D)$ for each n -disc $D \subset \subset \Omega$, where $U(\mathbf{x}) \geq 1$ and define an L^2 -function $W = U^\lambda$ for some $\lambda \in [1, 2)$. Suppose that*

$$\int_{\Omega} (a^{\alpha\beta} \partial_{\beta} W \partial_{\alpha} \zeta + f W \zeta) d\Omega \leq 0 \quad (87)$$

$\forall \zeta \in C_0^\infty(\Omega)$ with $\zeta(\mathbf{x}) \geq 0$, where the coefficients a and f are measurable with $f \in L^{n/2}(\Omega)$. Let the matrix a satisfy the ellipticity condition

$$C|\xi|^2 \leq a^{\alpha\beta}(\mathbf{x})\xi_{\alpha}\xi_{\beta}$$

for $|a(\mathbf{x})| \leq M$ at a.e. $\mathbf{x} \in \Omega$ and for all ξ . Moreover, let the growth condition (56) be satisfied. If $U \in L^2(\Omega)$, then U is bounded on each n -disc $D \subset \subset \Omega$ and satisfies

$$|U(\mathbf{x})|^2 \leq C a_0^{-n} \int_{D_{(R+a_0)}(\mathbf{x}_0)} |U(\mathbf{y})|^2 d\mathbf{y},$$

where $\mathbf{x}$ is a point of $D_R(\mathbf{x}_0) \subset \Omega$.

Slightly modified conditions will extend Morrey's result to $n = 2$; see [25], Sec. 5.4.

We apply Theorem A.1, taking $U(x) = H(Q(\mathbf{x})) + 1$ and $\lambda = 1$. Let a be given by the matrix α of Lemma 4 and let f be given by (55). Then f satisfies the local growth condition (56) by smoothness. The inequality of Lemma 3 completes the proof of Theorem 5.

7.2 Proof of Theorem 6

Without loss of generality, take p to be the origin of coordinates. Write the operator $\mathcal{L}_\omega$ in the form

$$\mathcal{L}_\omega(H) = \Delta H + \nabla \cdot [T(H)]$$

for

$$T = (-1)^{3n} * \{\omega \wedge * [\rho'(Q) dQ \wedge \omega]\}.$$

Then

$$\nabla [T(H)] = \partial_k (a^{jk} \partial_j H)$$

by arguments analogous to those of Lemma 4. Moreover, Lemma 4 implies that

$$|\nabla H|^2 \leq (1 + a^{jk}) \partial_j H \cdot \partial_k H \leq C |\nabla H|^2. \quad (88)$$

If f is given by (55), then

$$\int_{\Omega} \{[\nabla H + T(H)] \cdot \nabla \zeta - C f H \zeta\} d\Omega \leq 0 \quad (89)$$

for all nonnegative test functions $\zeta \in C_0^\infty(\Omega)$ which vanish in a neighborhood of the origin. Define

$$\zeta = (\eta \bar{\eta}_\nu)^2 H,$$

where $\eta \in C_0^\infty(D)$ for an n -disc D of radius R , containing the origin, and itself completely contained in the interior of Ω . Recalling that we have taken $H(0) = 0$ and that (9) implies that $H'(Q) > 0$, we conclude that H is a nonnegative function. Let $\bar{\eta}_\nu$ be given by the sequence [12]

$$\bar{\eta}_\nu(x) = \begin{cases} 0 & |x| \leq \nu^{-2} \\ \log(\nu^2 |x|) / \log(\nu^2 R) & \nu^{-2} < |x| < R \\ 1 & R \leq |x|. \end{cases}$$

Notice that $\bar{\eta}_\nu$ vanishes on a neighborhood of the origin for any finite parameter ν ; but as ν tends to infinity, $\bar{\eta}_\nu$ converges pointwise to 1, whereas $\nabla \bar{\eta}_\nu$ converges to zero in $L^n(D)$.

Inequality (89) now assumes the form

$$\begin{aligned} 0 &\leq \int_D [(\mathcal{L} + f) H] \cdot (\eta \bar{\eta}_\nu)^2 H * 1 \\ &= \int_D \{[\partial_k (1 + a^{jk}) \partial_j + f] H\} (\eta \bar{\eta}_\nu)^2 H * 1 = \\ &- \int (1 + a^{jk}) \partial_j H \cdot \partial_k [(\eta \bar{\eta}_\nu)^2 H] * 1 + \int_D f H^2 (\eta \bar{\eta}_\nu)^2 * 1, \end{aligned}$$

where

$$- \int_D (1 + a^{jk}) \partial_j H \cdot \partial_k [(\eta \bar{\eta}_\nu)^2 H] * 1 =$$

$$\begin{aligned}
& -2 \int_D (1 + a^{jk}) \bar{\eta}_\nu^2 \eta (\partial_k \eta) H \cdot \partial_j H * 1 \\
& -2 \int_D (1 + a^{jk}) \eta^2 \bar{\eta}_\nu (\partial_k \bar{\eta}_\nu) H \cdot \partial_j H * 1 \\
& - \int_D (1 + a^{jk}) (\eta \bar{\eta}_\nu)^2 (\partial_j H \cdot \partial_k H) * 1 \\
& \equiv - (2i_1 + 2i_2 + i_3).
\end{aligned}$$

That is,

$$\begin{aligned}
i_3 &= \int_D (1 + a^{jk}) (\eta \bar{\eta}_\nu)^2 (\partial_j H \cdot \partial_k H) * 1 \\
&\leq 2(|i_1| + |i_2|) + \int_D (\eta \bar{\eta}_\nu)^2 f H^2 * 1.
\end{aligned}$$

Estimating the integrals on the right individually, we have

$$\begin{aligned}
2i_1 &= 2 \int_D (1 + a^{jk}) \bar{\eta}_\nu^2 \eta (\partial_k \eta) H \cdot \partial_j H * 1 \leq \\
&\quad \varepsilon \int_D (\bar{\eta}_\nu \eta)^2 |\nabla H|^2 * 1 + \\
&\quad \frac{1}{\varepsilon} \int_D \bar{\eta}_\nu^2 |\nabla \eta|^2 H^2 * 1. \\
2i_2 &= 2 \int_D (1 + a^{jk}) \eta^2 \bar{\eta}_\nu (\partial_k \bar{\eta}_\nu) H \cdot \partial_j H * 1 \leq \\
&\quad \varepsilon \int_D (\bar{\eta}_\nu \eta)^2 |\nabla H|^2 * 1 + \frac{1}{\varepsilon} \int_D \eta^2 |\nabla \bar{\eta}_\nu|^2 H^2 * 1 \\
&\quad \equiv \varepsilon i_{21} + \frac{1}{\varepsilon} i_{22},
\end{aligned}$$

where the small constants ε depend on the constant C in the upper inequality of (88), and

$$i_{22} = \int_D \eta^2 |\nabla \bar{\eta}_\nu|^2 H^2 * 1 \leq C \|\nabla \bar{\eta}_\nu\|_n^2 \|\eta H\|_{2n/(n-2)}^2. \quad (90)$$

The right-hand side of (90) tends to zero as ν tends to infinity. Absorbing small constants on the left, now using the lower inequality of (88), we conclude that

$$\begin{aligned}
& (1 - 2\varepsilon) \int_D (\bar{\eta}_\nu \eta)^2 |\nabla H|^2 * 1 \leq \\
& C(\varepsilon) \left(\int_D \bar{\eta}_\nu^2 |\nabla \eta|^2 H^2 * 1 + \|\nabla \bar{\eta}_\nu\|_n^2 \|\eta H\|_{2n/(n-2)}^2 + \|f\|_{n/2} \|\eta \bar{\eta}_\nu H\|_{2n/(n-2)}^2 \right), \quad (91)
\end{aligned}$$

where

$$\begin{aligned} \|\eta \bar{\eta}_\nu H\|_{2n/(n-2)}^2 &\leq C \int_D |\nabla (\eta \bar{\eta}_\nu H)|^2 * 1 \leq \\ C \left(\int_D |\nabla (\eta \bar{\eta}_\nu)|^2 H^2 * 1 + \int_D (\eta \bar{\eta}_\nu)^2 |\nabla H|^2 * 1 \right). \end{aligned} \quad (92)$$

The first integral on the extreme right-hand side of inequality (92) has essentially already been estimated in (90), and the second can be subtracted from the left-hand side of (91) provided that its coefficient in (91), the $L^{n/2}$ -norm of f over Ω , is sufficiently small. Letting ν tend to infinity, we have in the limit

$$\int_D \eta^2 |\nabla H|^2 * 1 \leq C \int_D |\nabla \eta|^2 H^2 * 1.$$

We conclude that H is a weak solution in a neighborhood of the singularity. This completes the proof of Theorem 6.

7.3 Proof of Theorem 8

1. *Outline:* The result follows from the application of Lemmas 3 and 4 and Theorem 5, above, to the proofs of Theorem 6 and Corollary 7 of [27]. The absence of condition (33) results in a small change of L^p conditions on the solution.

2. *Details:* Initially proceed as Sec. 7.2, but choose the test of functions $\bar{\eta}_\nu$ to be a sequence of functions possessing the following properties:

- (a) $\bar{\eta}_\nu \in [0, 1] \forall \nu$;
- (b) $\bar{\eta}_\nu = 0$ in a neighborhood of $\Sigma \forall \nu$;
- (c) $\lim_{\nu \rightarrow \infty} \bar{\eta}_\nu = 1$ a.e.,
- (d) $\lim_{\nu \rightarrow \infty} \|\nabla \bar{\eta}_\nu\|_{L^{n-m-\varepsilon}} = 0$

(c.f. [33], Lemma 2 and p. 73). The function $\mathcal{F}$ is given by [32]

$$\mathcal{F}(H) = \begin{cases} H^{\gamma_2}, & 0 \leq H \leq \ell, \\ (1/\gamma_1) [\gamma_2 \ell^{\gamma_2-\gamma_1} H^{\gamma_1} - (\gamma_1 - \gamma_2) \ell^{\gamma_2}], & H \geq \ell. \end{cases}$$

The functions $\mathcal{F}(H)$ and

$$G(H) \equiv \mathcal{F}(H) F'(H) - \gamma_2$$

satisfy ([32], p. 280; see also Sec. 3 of [12]):

$$\mathcal{F}(H) \leq \frac{\gamma_2}{\gamma_1} \ell^{\gamma_2-\gamma_1} H^{\gamma_1};$$

$$H \mathcal{F}'(H) \leq \gamma_2 \mathcal{F};$$

$$\begin{aligned} |G(H)| &\leq \mathcal{F}(H)\mathcal{F}'(H); \\ G'(H) &\geq C\mathcal{F}'(H)^2. \end{aligned}$$

Replace the test function in Sec. 7.2 by the test function

$$\zeta = (\eta\bar{\eta}_\nu)^2 G(H),$$

where η is defined as in Sec. 7.2, and substitute this value into (89). We obtain

$$\begin{aligned} 0 &\leq \int_D \{ [\partial_k (1 + a^{jk}) \partial_j + f] H \} (\eta\bar{\eta}_\nu)^2 G(H) * 1 = \\ &- \int_D (1 + a^{jk}) \partial_j H \cdot \partial_k [(\eta\bar{\eta}_\nu)^2 G(H)] * 1 + \int_D f \eta^2 \bar{\eta}_\nu^2 H \cdot G(H) * 1, \end{aligned}$$

or

$$\begin{aligned} &2 \int_D (1 + a^{jk}) (\eta\bar{\eta}_\nu) \partial_k (\eta\bar{\eta}_\nu) (\partial_j H) G(H) * 1 + \\ &\int_D (1 + a^{jk}) (\eta\bar{\eta}_\nu)^2 G'(H) \partial_j H \partial_k H * 1 \leq \int_D f (\eta\bar{\eta}_\nu)^2 H \cdot G(H) * 1. \end{aligned}$$

Writing this inequality in the short-hand form

$$I_1 + I_2 \leq I_3, \quad (93)$$

we proceed analogously to inequalities (28)-(35) of [26], making the following changes from the notation of [26] to our notation: $\psi \rightarrow \bar{\eta}$, $Q \rightarrow H$, $H \rightarrow \mathcal{F}$, $\Xi \rightarrow G$, $\Phi \rightarrow f$. Explicitly,

$$\begin{aligned} I_1 &= 2 \int_D (1 + a^{jk}) (\eta\bar{\eta}_\nu) \partial_k (\eta\bar{\eta}_\nu) \cdot G(H) \partial_j H * 1 \\ &\geq -C \int_D (\eta\bar{\eta}_\nu) |\nabla (\eta\bar{\eta}_\nu)| |\mathcal{F}(H) \nabla \mathcal{F}(H)| * 1 \\ &\geq -\varepsilon \int_D (\eta\bar{\eta}_\nu)^2 |\nabla \mathcal{F}(H)|^2 * 1 - C(\varepsilon) \int_D |\nabla (\eta\bar{\eta})|^2 \mathcal{F}(H)^2 * 1; \\ I_2 &\geq C \int_D (\eta\bar{\eta}_\nu)^2 |\mathcal{F}'(H)|^2 |\nabla H|^2 * 1 \\ &\geq C \int_D (\eta\bar{\eta}_\nu)^2 |\nabla \mathcal{F}(H)|^2 * 1; \\ I_3 &= \int_D f (\eta\bar{\eta}_\nu)^2 H \cdot G(H) * 1 \leq \\ &\int_D |f| (\eta\bar{\eta}_\nu)^2 |H \cdot \mathcal{F}'(H)| |\mathcal{F}(H)| * 1 \leq \\ \gamma_2 \int_D |f| (\eta\bar{\eta}_\nu)^2 |\mathcal{F}(H)|^2 * 1 &\leq \|f\|_{n/2} \left(\int_D |\eta\bar{\eta}_\nu \mathcal{F}(H)|^{2n/(n-2)} * 1 \right)^{(n-2)/n}. \end{aligned} \quad (94)$$

As in (92), we apply the Sobolev inequality to the right-hand side of this expression, followed by the Minkowski and Schwartz inequalities:

$$\begin{aligned} & \left(\int_D |\eta \bar{\eta}_\nu \mathcal{F}(H)|^{2n/(n-2)} * 1 \right)^{(n-2)/n} \leq C \int_D |\nabla [\eta \bar{\eta}_\nu \mathcal{F}(H)]|^2 * 1 \\ & \leq C \left[\int_D |(\nabla \eta) \bar{\eta}_\nu \mathcal{F}(H)|^2 * 1 + \int_D |(\nabla \bar{\eta}_\nu) \eta \mathcal{F}(H)|^2 * 1 + \int_D (\eta \bar{\eta}_\nu)^2 |\nabla \mathcal{F}(H)|^2 * 1 \right] \\ & \equiv I_{31} + I_{32} + I_{33}. \end{aligned}$$

The term I_{33} can be subtracted from the left-hand-side of inequality (93), as its coefficient in (94), the $L^{n/2}$ -norm of f , is small on small discs as a consequence of condition (56).

$$\begin{aligned} I_{32} &= \int_D |(\nabla \bar{\eta}_\nu) \eta \mathcal{F}(H)|^2 * 1 \leq C(\gamma_1, \gamma_2, \ell) \int_D |(\nabla \bar{\eta}_\nu)^2 \eta^2 H^{2\gamma_1} * 1 \\ &\leq C \|\nabla \bar{\eta}_\nu\|_{n-m-\varepsilon}^2 \|H^{2\gamma_1}\|_\beta. \end{aligned}$$

Letting ν tend to infinity, the term on the right-hand side is zero for every value of ℓ . Now letting ℓ tend to infinity and using Fatou's inequality, we conclude that

$$\int_D \eta^2 |\nabla (H^{\gamma_2})|^2 * 1 \leq C \int_D |\nabla \eta|^2 H^{2\gamma_2} * 1.$$

Apply Theorem A.1, taking $U = H^{\gamma_2}$. Then $U \in H^{1,2}(D)$ and $W = U^\lambda$ satisfies inequality (87) for $\lambda = 1/\gamma_2$. Because $\gamma_2 > 1/2$, we can conclude that $\lambda < 2$ as required by Theorem A.1. We now want to check that we can choose $\gamma_2 \leq 1$, in order to obtain $\lambda \geq 1$ as also required by Theorem A.1. Because H lies in the space $L^{2\beta\gamma_1}(D) \cap L^{2\gamma_2}(D)$, we let $\gamma_1\beta = \gamma_2$. Substituting the definition of β , we find that we can choose $\gamma_2 \leq 1$ for $\gamma_1 > 1/2$ provided $m+4 < n$, which is satisfied by hypothesis. This completes the proof of Theorem 8.

8 Acknowledgments

We are grateful to Richard M. Schoen for suggesting reference [41], Marshall Slemrod for discussing the motivating example of Sec. 1.2, and Yisong Yang for discussing [8].

References

- [1] L. J. Alías, B. Palmer, A duality result between the minimal surface equation and the maximal surface equation, *An. Acad. Brasil. Ciênc.* **73** (2001) 161-164.
- [2] H. Beirão de Veiga, On non-Newtonian p -fluids. The pseudo-plastic case, *J. Math. Anal. Appl.* **344** (2008), 175-185.

- [3] L. Bers, Results and conjectures in the mathematical theory of subsonic and transonic gas flows, *Commun. Pure Appl. Math.* **7** (1954), 79-109.
- [4] L. Bers, *Mathematical Aspects of Subsonic and Transonic Gas Dynamics*, Wiley, New York, 1958.
- [5] O. I. Bogoyavlenskii, Integrable Euler equations on Lie algebras arising in problems of mathematical physics [in Russian], *Izvestiya Akad. Nauk SSSR* **48**, No. 5 (1984); translation in *Mathematics of the USSR-Izvestiya* **25**, No. 2 (1985), 207-257.
- [6] M. Born and L. Infeld, Foundation of a new field theory, *Proc. R. Soc. London Ser. A* **144** (1934) 425-451.
- [7] E. Calabi, Examples of Bernstein problems for some nonlinear equations, *Proceedings of the Symposium on Global Analysis*, University of California at Berkeley, 1968, pp. 223-230.
- [8] E. DiBenedetto, $C^{1+\alpha}$ local regularity of weak solutions to degenerate elliptic equations, *Nonlinear Analysis T. M. A.* **7**, No. 8 (1983), 827-850.
- [9] H. Dinh and G. F. Carey, Some results concerning approximation of regularized compressible flow, *Int. J. Num. Meth. Fluids* **5** (1985), 299-302.
- [10] D. G. B. Edelen, *Applied Exterior Calculus*, Wiley, New York, 1985.
- [11] G. W. Gibbons, Born-Infeld particles and Dirichlet p -branes, *Nucl. Phys. B* **514**, 603-639.
- [12] B. Gidas and J. Spruck, Global and local behavior of positive solutions of nonlinear elliptic equations, *Commun. Pure Appl. Math.* **4** (1981), 525-598.
- [13] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin, 1983.
- [14] H. Helmholtz, Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen, *J. Reine Angew. Math.* **55** (1858), 25-55.
- [15] T. Isobe, A regularity result for a class of degenerate Yang-Mills connections in critical dimensions, *Forum Math.* **20** (2008), 1109-1139.
- [16] T. Iwaniec, C. Scott, and B. Stroffolini, Nonlinear Hodge theory on manifolds with boundary, *Annali Mat. Pura Appl.* **177** (1999) 37-115.
- [17] E. Kreyszig, On the theory of minimal surfaces, in: Th. M. Rassias, ed., *The Problem of Plateau: A Tribute to Jesse Douglas and Tibor Radó*, World Scientific, Singapore, 1992, pp. 138-164.
- [18] O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and Quasilinear Elliptic Equations*, Academic Press, New York, 1968.

- [19] R. Magnanini and G. Talenti, On complex-valued solutions to a 2D eikonal equation. Part one: qualitative properties, *Contemporary Math.* **283** (1999) 203-229.
- [20] R. Magnanini and G. Talenti, On complex-valued solutions to a two-dimensional eikonal equation. II. Existence theorems, *SIAM J. Math. Anal.* **34** (2003), no. 4, 805-835.
- [21] R. Magnanini and G. Talenti, On complex-valued solutions to a 2D eikonal equation. III. Analysis of a *Bäcklund* transformation, *Appl. Anal.* **85** (2006), no. 1-3, 249-276.
- [22] K. B. Marathe and G. Martucci, *The Mathematical Foundations of Gauge Theories*, North-Holland, Amsterdam, 1992.
- [23] A. Marini, The generalized Neumann problem for Yang-Mills connections, *Commun. Partial Differential Equations* **24** (1999), 665-681.
- [24] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, Freeman, New York, 1973.
- [25] C. B. Morrey, *Multiple Integrals in the Calculus of Variations*, Springer-Verlag, Berlin, 1966.
- [26] T. H. Otway, Nonlinear Hodge maps, *J. Math. Phys.* **41** (2000), 5745-5766.
- [27] T. H. Otway, Maps and fields with compressible density, *Rendiconti del Seminario Matematico dell' Università di Padova* **111** (2004), 133-159.
- [28] T. H. Otway, Variational equations on mixed Riemannian-Lorentzian metrics, *J. Geometry & Phys.* **58** (2008), 1043-1061.
- [29] E. Poisson, *A Relativist's Toolkit: The Mathematics of Black-Hole Mechanics*, Cambridge, 2004.
- [30] D. Riabouchinsky, Sur l'analogie hydraulique des mouvements d'un fluide compressible, *Comptes Rendus Academie des Sciences, Paris* **195** (1932), 998.
- [31] C. Rogers and W. K. Schief, *Bäcklund and Darboux Transformations, Geometry and Modern Applications of Soliton Theory*, Cambridge University Press, Cambridge, 2002.
- [32] J. Serrin, Local behavior of solutions of quasilinear equations, *Acta Math.* **111** (1964), 247-302.
- [33] J. Serrin, Removable singularities of solutions of elliptic equations, *Arch. Ration. Mech. Anal.* **17** (1964), 67-78.
- [34] M. Shiffman, On the existence of subsonic flows of a compressible fluid, *J. Rat. Mech. Anal.* **1** (1952), 605-652.

- [35] L. M. Sibner, An existence theorem for a non-regular variational problem, *Manuscripta Math.* **43**, 45-72 (1983).
- [36] L. M. Sibner and R. J. Sibner, A nonlinear Hodge-de Rham theorem, *Acta Math.* **125** (1970), 57-73.
- [37] L. M. Sibner and R. J. Sibner, Nonlinear Hodge theory: Applications, *Advances in Math.* **31** (1979), 1-15.
- [38] L. M. Sibner, R. J. Sibner, and Y. Yang, Generalized Bernstein property and gravitational strings in Born-Infeld theory, *Nonlinearity* **20** (2007) 1193-1213.
- [39] P. D. Smith, Nonlinear Hodge theory on punctured Riemannian manifolds, *Indiana Univ. Math. J.* **31** (1982), 553-577.
- [40] J. J. Stoker, *Water Waves*, Interscience, New York, 1987.
- [41] B. Tabak, A geometric characterization of harmonic diffeomorphisms between surfaces, *Math. Ann.* **270** (1985), 147-157.
- [42] K. K. Uhlenbeck, Regularity for a class of nonlinear elliptic systems, *Acta Math.* **138** (1977), 219-240.
- [43] Y. Yang, Classical solutions in the Born-Infeld theory, *Proc. R. Soc. Lond. Ser. A* **456** (2000) 615-640.